

Composition followed by differentiation between weighted Bergman spaces and weighted Banach spaces of holomorphic functions

Elke Wolf

Abstract. Let ϕ be an analytic self-map of the open unit disk \mathbb{D} in the complex plane. Such a map induces through composition a linear composition operator $C_\phi : f \mapsto f \circ \phi$. We are interested in the combination of C_ϕ with the differentiation operator D , that is in the operator $DC_\phi : f \mapsto \phi' \cdot (f \circ \phi)$ acting between weighted Bergman spaces and weighted Banach spaces of holomorphic functions.

Mathematics subject classification: 47B33, 47B38.

Keywords and phrases: Composition operator, differentiation operator, weighted Bergman spaces, weighted Banach spaces of holomorphic functions.

1 Introduction

Let \mathbb{D} denote the open unit disk in the complex plane. For an analytic self-map ϕ of \mathbb{D} the classical *composition operator* C_ϕ is given by

$$C_\phi : H(\mathbb{D}) \rightarrow H(\mathbb{D}), f \mapsto f \circ \phi,$$

where $H(\mathbb{D})$ denotes the set of all analytic functions on \mathbb{D} . Combining this with differentiation we obtain the operator

$$DC_\phi : H(\mathbb{D}) \rightarrow H(\mathbb{D}), f \mapsto \phi' \cdot (f \circ \phi).$$

Composition operators occur naturally in various problems and therefore have been widely investigated. An overview of results in the classical setting of the Hardy spaces as well as an introduction to composition operators is given in the excellent monographs by Cowen and MacCluer [5] and Shapiro [8].

Next, let us explain the setting in which we are interested. Bounded and continuous functions $v : \mathbb{D} \rightarrow]0, \infty[$ are called *weights*. For such a weight v we define

$$H_v^\infty := \{f \in H(\mathbb{D}); \|f\|_v := \sup_{z \in \mathbb{D}} v(z)|f(z)| < \infty\}.$$

Since, endowed with the weighted sup-norm $\|\cdot\|_v$, this is a Banach space, we say that H_v^∞ is a *weighted Banach space of holomorphic functions*. These spaces arise naturally in several problems related to e. g. complex analysis, spectral theory, Fourier analysis, partial differential and convolution equations. Concrete examples may be

found in [3]. Weighted Banach spaces of holomorphic functions have been studied deeply in [2], but also in [4] and [1].

The *weighted Bergman space* is defined to be the collection of all analytic functions $f \in H(\mathbb{D})$ such that

$$A_{v,p} := \left\{ f \in H(\mathbb{D}); \|f\|_{v,p} := \left(\int_{\mathbb{D}} |f(z)|^p v(z) dA(z) \right)^{\frac{1}{p}} < \infty \right\}, 1 \leq p < \infty$$

where $dA(z)$ denotes the normalized area measure. The investigation of Bergman spaces has quite a long and rich history. An excellent introduction to Bergman spaces is given in [6].

In this article we characterize boundedness and compactness of operators $DC_\phi : A_{v,p} \rightarrow H_w^\infty$ in terms of the involved self-map ϕ and the weights v and w .

2 Basics

We study weighted spaces generated by the following class of weights. Let ν be a holomorphic function on \mathbb{D} that does not vanish and is strictly positive on $[0, 1[$. Moreover, we assume that $\lim_{r \rightarrow 1} \nu(r) = 0$. Then we define the weight v in the following way

$$v(z) := \nu(|z|^2) \text{ for every } z \in \mathbb{D}. \quad (1)$$

Examples include all the famous and popular weights, such as

1. the *standard weights* $v(z) = (1 - |z|^2)^\alpha$, $\alpha \geq 1$,
2. the *logarithmic weights* $v(z) = (1 - \log(1 - |z|^2))^\beta$, $\beta > 0$.
3. the *exponential weights* $v(z) = e^{-\frac{1}{(1-|z|^2)^\alpha}}$, $\alpha \geq 1$.

For a fixed point $a \in \mathbb{D}$, we introduce a function

$$v_a(z) := \nu(\bar{a}z) \text{ for every } z \in \mathbb{D}.$$

Since ν is holomorphic on \mathbb{D} , so is the function v_a . Moreover, in particular, we will often assume that there is a constant $C > 0$ such that

$$\sup_{a \in \mathbb{D}} \sup_{z \in \mathbb{D}} \frac{v(z)}{|v_a(z)|} \leq C. \quad (2)$$

In the sequel we analyze which role condition (2) plays in the zoo of conditions on weights. Lusky [7] studied weights satisfying the following conditions (L1) and (L2) (renamed after the author) which are defined as follows

$$(L1) \inf_{n \in \mathbb{N}} \frac{v(1 - 2^{-n-1})}{v(1 - 2^{-n})} > 0 \text{ and } (L2) \limsup_{n \rightarrow \infty} \frac{v(1 - 2^{-n-j})}{v(1 - 2^{-n})} < 1 \text{ for some } j \in \mathbb{N}.$$

Actually, weights which enjoy both conditions (L1) and (L2) are normal weights in the sense of Shields and Williams (see [9]). Obviously condition (2) is connected with condition (L2) in the following way. If we change (2) as follows

$$\sup_{a \in \mathbb{D}} \sup_{z \in \mathbb{D}} \frac{v(z)}{|v_a(z)|} < 1, \quad (3)$$

then (L2) is equivalent with (3) if we assume that $|\nu(z)| \geq \nu(|z|)$ for every $z \in \mathbb{D}$. To show this, let us first assume that (L2) holds. Hence we can find $j \in \mathbb{N}$ such that

$$\frac{v(1 - 2^{-n-j})}{v(1 - 2^{-n})} < 1 \text{ for every } n \in \mathbb{N}.$$

Next, we fix $z \in \mathbb{D}$ and $a \in \mathbb{D}$. Then we can find $n \in \mathbb{N}$ such that

$$|z| \geq 1 - 2^{-n-j} \text{ and } |az| < 1 - 2^{-n}.$$

Now,

$$\frac{v(z)}{|\nu(az)|} \leq \frac{v(1 - 2^{-n-j})}{v(1 - 2^{-n})} < 1 \text{ for every } n \in \mathbb{N}.$$

Conversely, we suppose that (3) is satisfied. We take $j = 1$, fix $n \in \mathbb{N}$ and select

$$a_n := \frac{(1 - 2^{-n})^2}{(1 - 2^{-n-1})}.$$

We obtain

$$\frac{v(1 - 2^{-n-1})}{v(1 - 2^{-n})} \leq \frac{v(z)}{|\nu(az)|} \leq \sup_{a \in \mathbb{D}} \sup_{z \in \mathbb{D}} \frac{v(z)}{|v_a(z)|} < 1.$$

Thus, under some additional assumptions (2) is a weaker version of (L2). Calculations show that the standard weights as well as the logarithmic weights satisfy condition (2), while the exponential weights do not fulfill condition (2).

Finally, we need some geometric data of the unit disk. A very important tool when dealing with operators such as defined above is the so-called *pseudohyperbolic metric* given by

$$\rho(z, a) := |\sigma_a(z)|,$$

where $\sigma_a(z) = \frac{a-z}{1-\bar{a}z}$, $z, a \in \mathbb{D}$, is the Möbius transformation which interchanges a and 0.

3 Results

Lemma 1. *Let $v(z) = f(|z|)$ for every $z \in \mathbb{D}$, where $f \in H(\mathbb{D})$ is a function whose Taylor series (at 0) has nonnegative coefficients. We assume additionally that v satisfies condition (2). Then there is a constant $C > 0$ such that*

$$|f(z)| \leq C^{\frac{1}{p}} \frac{\|f\|_{v,p}}{v(0)^{\frac{1}{p}} (1 - |z|^2)^{\frac{2}{p}} v(z)^{\frac{1}{p}}}.$$

Proof. Recall that a weight v as defined above may be written as

$$v(z) := \max\{|g(\lambda z)|; |\lambda| = 1\} \text{ for every } z \in \mathbb{D}.$$

We will write $g_\lambda(z) := g(\lambda z)$ for every $z \in \mathbb{D}$. Next, fix $\lambda \in \mathbb{C}$ with $|\lambda| = 1$. Moreover, let $\alpha \in \mathbb{D}$ be an arbitrary point. We consider the map

$$T_{\alpha,\lambda} : A_v^p \rightarrow A_v^p, \quad T_{\alpha,\lambda}f(z) = f(\sigma_\alpha(z))\sigma'_\alpha(z)^{\frac{2}{p}}g_\lambda(\sigma_\alpha(z))^{\frac{1}{p}}.$$

Then a change of variables yields

$$\begin{aligned} \|T_{\alpha,\lambda}f\|_{v,p}^p &= \int_{\mathbb{D}} v(z)|f(\sigma_\alpha(z))|^p |\sigma'_\alpha(z)|^2 |g_\lambda(\sigma_\alpha(z))| dA(z) \\ &\leq \int_{\mathbb{D}} |f(\sigma_\alpha(z))| \frac{v(z)}{v(\sigma_\alpha(z))} |\sigma'_\alpha(z)| dA(z) \\ &\leq C \int_{\mathbb{D}} |f(\sigma_\alpha(z))| v(\sigma_\alpha(z)) |\sigma'_\alpha(z)|^2 dA(z) \\ &\leq C \int_{\mathbb{D}} v(t)|f(t)|^p dA(t) = C\|f\|_{v,p}^p. \end{aligned}$$

Now put $h_\lambda(z) := T_{\alpha,\lambda}(z)$ for every $z \in \mathbb{D}$. By the mean value property we obtain

$$v(0)|h_\lambda(0)|^p \leq \int_{\mathbb{D}} v(z)|h_\lambda(z)|^p dA(z) = \|h_\lambda\|_{v,p}^p \leq C\|f\|_{v,p}^p.$$

Hence

$$v(0)|h_\lambda(0)|^p = v(0)|f(\alpha)|^p (1 - |\alpha|^2)^2 |g_\lambda(\alpha)| \leq C\|f\|_{v,p}^p.$$

Since λ was arbitrary we obtain that

$$v(0)|f(\alpha)|^p (1 - |\alpha|^2)^2 v(\alpha) \leq C\|f\|_{v,p}^p$$

Thus,

$$|f(\alpha)| \leq C^{\frac{1}{p}} \frac{\|f\|_{v,p}}{v(0)^{\frac{1}{p}} (1 - |\alpha|^2)^{\frac{2}{p}} v(\alpha)^{\frac{1}{p}}}.$$

Since α was arbitrary, the claim follows. \square

Lemma 2. *Let $v(z) = f(|z|)$ for every $z \in \mathbb{D}$, where $f \in H(\mathbb{D})$ is a function whose Taylor series (at 0) has nonnegative coefficients. We assume additionally that v satisfies condition (2). Then for every $f \in A_v^p$ there is $C_v > 0$ such that*

$$|f(z) - f(w)| \leq C_v \|f\|_{v,p} \max \left\{ \frac{1}{(1 - |z|^2)^{\frac{2}{p}} v(z)^{\frac{1}{p}}}, \frac{1}{(1 - |w|^2)^{\frac{2}{p}} v(w)^{\frac{1}{p}}} \right\} \rho(z, w)$$

for every $z, w \in \mathbb{D}$.

Proof. The proof is completely analogous to the proof given in [10]. Hence we omit it here. \square

Lemma 3. *Let $v(z) = f(|z|)$ for every $z \in \mathbb{D}$, where $f \in H(\mathbb{D})$ is a function whose Taylor series (at 0) has nonnegative coefficients. We assume additionally that v satisfies condition (2). Then for $f \in H_v^\infty$ and $z \in \mathbb{D}$:*

$$|f'(z)| \leq \frac{M}{v(0)^{\frac{1}{p}}(1-|z|^2)^{1+\frac{2}{p}}v(z)^{\frac{1}{p}}} \|f\|_{v,p}.$$

Proof. By Lemma 2 we have that

$$|f(z) - f(w)| \leq \frac{M}{v(0)^{\frac{1}{p}}} \max \left\{ \frac{1}{(1-|z|^2)^{\frac{2}{p}}v(z)^{\frac{1}{p}}}, \frac{1}{(1-|w|^2)^{\frac{2}{p}}v(w)^{\frac{1}{p}}} \right\} \rho(z, w) \|f\|_{v,p}.$$

Now

$$\begin{aligned} & \left| \frac{f(z+h) - f(z)}{|h|} \right| \leq \\ & \leq \frac{M}{v(0)^{\frac{1}{p}}|h|} \max \left\{ \frac{1}{(1-|z+h|^2)^{\frac{2}{p}}v(z+h)^{\frac{1}{p}}}, \frac{1}{(1-|z|^2)^{\frac{2}{p}}v(z)^{\frac{1}{p}}} \right\} \rho(z+h, z) \|f\|_{v,p} \\ & = \frac{M}{v(0)^{\frac{1}{p}}|h|} \max \left\{ \frac{1}{(1-|z+h|^2)^{\frac{2}{p}}v(z+h)^{\frac{1}{p}}}, \frac{1}{(1-|z|^2)^{\frac{2}{p}}v(z)^{\frac{1}{p}}} \right\} \left| \frac{z+h-z}{1-\overline{z+h}z} \right| \|f\|_{v,p} \\ & = \frac{M}{v(0)^{\frac{1}{p}}} \max \left\{ \frac{1}{(1-|z+h|^2)^{\frac{2}{p}}v(z+h)^{\frac{1}{p}}}, \frac{1}{(1-|z|^2)^{\frac{2}{p}}v(z)^{\frac{1}{p}}} \right\} \left| \frac{1}{1-\overline{z+h}z} \right| \|f\|_{v,p}. \end{aligned}$$

Finally, let h tend to zero and obtain

$$|f'(z)| \leq \frac{M}{v(0)^{\frac{1}{p}}(1-|z|^2)^{1+\frac{2}{p}}v(z)^{\frac{1}{p}}} \|f\|_{v,p}.$$

□

Proposition 1. *Let $v(z) = f(|z|)$ for every $z \in \mathbb{D}$, where $f \in H(\mathbb{D})$ is a function whose Taylor series (at 0) has nonnegative coefficients. We assume additionally that v satisfies condition (2). Then $DC_\phi : A_{v,p} \rightarrow H_w^\infty$ is bounded if and only if*

$$\sup_{z \in \mathbb{D}} \frac{w(z)|\phi'(z)|}{(1-|\phi(z)|^2)^{1+\frac{2}{p}}v(\phi(z))^{\frac{1}{p}}} < \infty. \quad (4)$$

Proof. First, we assume that (4) is satisfied. Applying Lemma 1 we obtain

$$\|DC_\phi f\|_w = \sup_{z \in \mathbb{D}} w(z)|\phi'(z)| |f'(\phi(z))| \leq C \sup_{z \in \mathbb{D}} \frac{w(z)|\phi'(z)|}{(1-|\phi(z)|^2)^{1+\frac{2}{p}}v(\phi(z))^{\frac{1}{p}}}.$$

Hence $DC_\phi : A_{v,p} \rightarrow H_w^\infty$ must be bounded.

Conversely, let $a \in \mathbb{D}$ be arbitrary. Then there exists f_a^p in the unit ball of H_v^∞ such that $|f_a(a)|^p = \frac{1}{v(a)}$. Now put

$$g_a(z) := f_a(z)\sigma_a(z) \text{ for every } z \in \mathbb{D}.$$

Then $\|g_a\|_{v,p}^p = \int_{\mathbb{D}} |g_a(z)|^p v(z) dA(z) \leq \sup_{z \in \mathbb{D}} v(z) |f_a(z)|^p \int_{\mathbb{D}} |\sigma(z)|^p dA(z) \leq K$.
 Moreover,

$$g'_a(z) = f'_a(z)\sigma_a(z) + f_a(z)\sigma'_a(z) \text{ for every } z \in \mathbb{D}.$$

Next, we assume that there is a sequence $(z_n)_n \subset \mathbb{D}$ such that $|\phi(z_n)| \rightarrow 1$ and

$$\frac{w(z_n)|\phi'(z_n)|}{(1 - |\phi(z_n)|^2)^{1+\frac{2}{p}} v(\phi(z_n))^{\frac{1}{p}}} \geq n \text{ for every } n \in \mathbb{N}.$$

Thus consider now $g_n(z) := g_{\phi(z_n)}(z)$ for every $n \in \mathbb{N}$ as defined above. Obviously $(g_n)_n$ is contained in the closed unit ball of $A_{v,p}$ and

$$c \geq w(z_n)|\phi'(z_n)||g'_n(\phi(z_n))| = \frac{w(z_n)|\phi'(z_n)|}{v(\phi(z_n))^{\frac{1}{p}}(1 - |\phi(z_n)|^2)^{1+\frac{2}{p}}} \geq n$$

for every $n \in \mathbb{N}$ which is a contradiction. \square

Proposition 2. *Let $v(z) = f(|z|)$, $z \in \mathbb{D}$, where $f \in H(\mathbb{D})$ is a function whose Taylor series (at 0) has nonnegative coefficients. Moreover, we assume that v satisfies (2). Then the operator $DC_\phi : A_{v,p} \rightarrow H_w^\infty$ is compact if and only if*

$$\limsup_{|\phi(z)| \rightarrow 1} \frac{w(z)|\phi'(z)|}{(1 - |\phi(z)|^2)^{1+\frac{2}{p}} v(\phi(z))^{\frac{1}{p}}}.$$

Proof. Let $(f_n)_n$ be a bounded sequence in $A_{v,p}$ that converges to zero uniformly on the compact subsets of \mathbb{D} . Let $M := \sup_n \|f_n\|_{v,p} < \infty$. Given $\varepsilon > 0$ there is $r > 0$ such that if $|\phi(z)| > 0$, then

$$\frac{w(z)|\phi'(z)|}{(1 - |\phi(z)|^2)^{1+\frac{2}{p}} v(\phi(z))^{\frac{1}{p}}} \leq \frac{\varepsilon}{2C_v}.$$

On the other hand, since $f_n \rightarrow 0$ uniformly on $\{u; |u| \leq r\}$, there is an $n_0 \in \mathbb{N}$ such that if $|\phi(z)| \leq r$ and $n \geq n_0$, then $w(z)|f'_n(\phi(z))||\phi'(z)| < \frac{\varepsilon}{2}$. Now, an application of Lemma 3 yields

$$\begin{aligned} \sup_{z \in \mathbb{D}} w(z)|f'_n(\phi(z))||\phi'(z)| &\leq \sup_{|\phi(z)| \leq r} w(z)|f'_n(\phi(z))||\phi'(z)| + \\ &+ \sup_{|\phi(z)| > r} w(z)|f'_n(\phi(z))||\phi'(z)| \leq \frac{\varepsilon}{2} + \sup_{|\phi(z)| > r} \frac{C_v w(z)|\phi'(z)|}{(1 - |\phi(z)|^2)^{\frac{2}{p}+1} v(\phi(z))^{\frac{1}{p}}} < \varepsilon. \end{aligned}$$

Thus, the claim follows.

Conversely, we suppose that $DC_\phi : A_{v,p} \rightarrow H_w^\infty$ is compact and that there are $\delta > 0$ and $(z_n)_n \subset \mathbb{D}$ with $|\phi(z_n)| \rightarrow 1$ such that

$$\frac{w(z_n)|\phi'(z_n)|}{(1 - |\phi(z_n)|^2)^{1+\frac{2}{p}} v(\phi(z_n))^{\frac{1}{p}}} \geq \delta.$$

Since $|\phi(z_n)| \rightarrow 1$ there exist natural numbers $\alpha(n)$ with $\lim_{n \rightarrow \infty} \alpha(n) = \infty$ such that $|\phi(z_n)|^{\alpha(n)} \geq \frac{1}{2}$ for every $n \in \mathbb{N}$.

Next, for every $n \in \mathbb{N}$ we consider the function

$$g_n(z) := f_n(z) \sigma_{\phi(z_n)}^{1+\frac{2}{p}} z^{\alpha(n)},$$

where $f_n^p \in H_v^\infty$ such that $\|f_n^p\|_v \leq 1$ and $|f_n(\phi(z_n))|^p = \frac{1}{\tilde{v}(\phi(z_n))}$. Then we obtain

$$\begin{aligned} \|DC_\phi f_n\|_w &\geq w(z_n) |\phi'(z_n)| |f_n'(\phi(z_n))| \\ &\geq \frac{w(z_n) |\phi'(z_n)| |\phi(z_n)|^{\alpha(n)}}{\tilde{v}(\phi(z_n))^{\frac{1}{p}} (1 - |\phi(z_n)|^2)^{1+\frac{2}{p}}} \\ &\geq \frac{1}{2} \frac{w(z_n) |\phi'(z_n)|}{\tilde{v}(\phi(z_n))^{\frac{1}{p}} (1 - |\phi(z_n)|^2)^{1+\frac{2}{p}}} \geq \frac{1}{2} \delta. \end{aligned}$$

This is a contradiction. □

References

- [1] BIERSTEDT K. D., BONET J., GALBIS A. *Weighted spaces of holomorphic functions on balanced domains*. Michigan Math. J., 1993, **40**, No. 2, 271–297.
- [2] BIERSTEDT K. D., BONET J., TASKINEN J. *Associated weights and spaces of holomorphic functions*. Studia Math., 1998, **127**, No. 2, 137–168.
- [3] BIERSTEDT K. D., MEISE R., SUMMERS W. H. *A projective description of weighted inductive limits*. Trans. Am. Math. Soc., 1982, **272**, No. 1, 107–160.
- [4] BIERSTEDT K. D., SUMMERS W. H. *Biduals of weighted Banach spaces of holomorphic functions*. J. Austral. Math. Soc., Ser. A., 1993, **54**, 70–79.
- [5] COWEN C., MACCLUER B. *Composition Operators on Spaces of Analytic Functions*. CRC Press, Boca Raton, 1995.
- [6] DUREN P., SCHUSTER A. *Bergman spaces, Mathematical Surveys and Monographs 100*. Amer. Math. Soc, Providence, RI, 2004.
- [7] LUSKY W. *On the structure of $Hv_0(D)$ and $hv_0(D)$* . Math. Nachr., 1992, **159**, 279–289.
- [8] SHAPIRO J. H. *Composition Operators and Classical Function Theory*. Springer, 1993.
- [9] SHIELDS A. L., WILLIAMS D. L. *Bounded projections, duality and multipliers in spaces of harmonic functions*. J. Reine Angew. Math., 1978, 299/300, 256–279.
- [10] WOLF E. *Differences of composition operators between weighted Bergman spaces and weighted Banach spaces of holomorphic functions*. Glasgow Math. J., 2010, **52**, No. 2, 325–332.

ELKE WOLF
 Mathematical Institute
 University of Paderborn
 Warburger Str. 100
 33098 Paderborn
 Germany

E-mail: lichte@math.uni-paderborn.de

Received September 30, 2013