

## Sensitivity analysis of efficient solution in vector MINMAX boolean programming problem \*

Vladimir A. Emelichev, Vitaly N. Krichko, Yury V. Nikulin

### Abstract

We consider a multiple criterion Boolean programming problem with MINMAX partial criteria. The extreme level of independent perturbations of partial criteria parameters such that efficient (Pareto optimal) solution preserves optimality was obtained.

**MSC: 90C29, 90C31**

**Key words and phrases:** vector MINMAX Boolean programming problem, efficient solution, stability radius.

Let  $C = (c_{ij}) \in \mathbf{R}^{n \times m}$ ,  $n, m \in \mathbf{N}$ ,  $m \geq 2$ ,  $C_i = (c_{i1}, c_{i2}, \dots, c_{im})$ ,  $\mathbf{E}^m = \{0, 1\}^m$ ,  $T$  be the non-empty subset of the permutations set  $S_m$  which is defined on the set  $N_m = \{1, 2, \dots, m\}$ . On the set of non-zero solutions (i.e. Boolean non-zero vectors)  $X \subseteq \mathbf{E}^m$ ,  $|X| > 1$ , we define the vector criterion

$$f(x, C) = (f_1(x, C_1), f_2(x, C_2), \dots, f_n(x, C_n)) \longrightarrow \min_{x \in X}.$$

The components (partial criteria) are functions

$$f_i(x, C_i) = \max_{t \in T} \sum_{j \in N(x)} c_{it(j)}, \quad i \in N_n,$$

where

$$t = \begin{bmatrix} 1 & 2 & \dots & m \\ t(1) & t(2) & \dots & t(m) \end{bmatrix}, \quad N(x) = \{j \in N_m : x_j = 1\}.$$

---

©2002 by V.A.Emelichev, V.N.Krichko, Yu.V.Nikulin

\*This work was partially supported by State program of Fundamental Investigations of Republic of Belarus "Mathematical structure" (grant 913/28).

Suppose  $C_i[t] = (c_{it(1)}, c_{it(2)}, \dots, c_{it(m)})$ . Then we can rewrite partial criteria in the following form

$$f_i(x, C_i) = \max_{t \in T} C_i[t]x, \quad i \in N_n,$$

where

$$x = (x_1, x_2, \dots, x_m)^T.$$

The problem of finding the set of *efficient solutions* (the Pareto set)

$$P^n(C) = \{x \in X \mid \pi(x, C) = \emptyset\}$$

we call a *vector minimax Boolean programming problem* and write  $Z^n(C)$ , where

$$\pi(x, C) = \{x' \in X : q(x, x', C) \geq 0_{(n)}, q(x, x', C) \neq 0_{(n)}\}.$$

$$q(x, x', C) = (q_1(x, x', C_1), q_2(x, x', C_2), \dots, q_n(x, x', C_n)),$$

$$q_i(x, x', C_i) = f_i(x, C_i) - f_i(x', C_i), \quad i \in N_n, \quad 0_{(n)} = (0, 0, \dots, 0) \in \mathbf{R}^n.$$

By analogy with [1 – 4], where the stability radius of efficient solution in different optimization problems was studied, the number

$$\rho^n(x^0, C) = \begin{cases} \sup \Omega & \text{if } \Omega \neq \emptyset, \\ 0 & \text{otherwise} \end{cases}$$

is called *the stability radius of the efficient solution*  $x^0 \in P^n(C)$ . Here

$$\Omega = \{\varepsilon > 0 : \forall C' \in \mathfrak{R}(\varepsilon) (x^0 \in P^n(C + C'))\},$$

$$\mathfrak{R}(\varepsilon) = \{C' \in \mathbf{R}^{n \times m} : \|C'\|_\infty < \varepsilon\},$$

$$\|C'\|_\infty = \max\{|c'_{ij}| : (i, j) \in N_n \times N_m\}, \quad C' = (c'_{ij}) \in \mathbf{R}^{n \times m}.$$

We consider  $\rho^n(x^0, C) = \infty$  if for any matrix  $C' \in \mathbf{R}^{n \times m}$

$$x' \in P^n(C + C').$$

For any  $x^0 \neq x$  and any permutation  $t \in T$  we introduce the following notations:

$$T(x^0, x) = \{t \in T : \forall t' \in T (N(x^0, t) \neq N(x, t'))\},$$

$$N(x, t) = \{t(j) : j \in N_m \ \& \ x_j = 1\},$$

$$\bar{T}(x^0, x) = T \setminus T(x^0, x).$$

**Lemma 1** Assume that  $x^0 \neq x$ ,  $x^0, x \in X$   $t^0 \in \bar{T}(x^0, x)$ . Then

$$C_i[t^0]x^0 \leq f_i(x, C_i)$$

for any index  $i \in N_n$  and matrix  $C \in \mathbf{R}^{n \times m}$ .

**Proof.** Let  $t^0 \in \bar{T}(x^0, x)$ . Then there exists  $t' \in T$  such that  $N(x^0, t^0) = N(x, t')$ . So for any  $i \in N_n$  we have

$$C_i[t^0]x^0 = C_i[t']x \leq \max_{t \in T} C_i[t]x = f_i(x, C_i).$$

Lemma 1 is proved.

The efficient solution  $x^0$  is called *trivial* if the set  $T(x^0, x)$  is empty for any  $x \in X \setminus \{x^0\}$  and *non-trivial* otherwise.

**Theorem 1** The stability radius  $\rho^n(x^0, C)$  of any trivial solution  $x^0$  of the problem  $Z^n(C)$  is infinite.

**Proof.** Let  $x^0 \in P^n(C)$ . Since  $x$  trivial, the equality  $T = \bar{T}(x^0, x)$  is true for any  $x \in X \setminus \{x^0\}$ . By lemma 1, the inequality

$$(C + C')_i[t^0]x^0 \leq f_i(x, C_i + C'_i)$$

holds for any  $x \in X \setminus \{x^0\}$ ,  $t^0 \in T$ ,  $i \in N_n$ ,  $C' \in \mathbf{R}^{n \times m}$ . Hence

$$q(x^0, x, C + C') \leq 0_{(n)}.$$

So the solution  $x^0 \in P^n(C)$  preserves the efficiency for any independent perturbations of matrix  $C$ . Thus  $\rho^n(x^0, C) = \infty$ . Theorem 1 is proved.

By definition, put

$$X(x^0) = \{x \in X \setminus \{x^0\} : T(x^0, x) \neq \emptyset\}.$$

**Lemma 2** Let  $x^0$  be non-trivial efficient solution of the problem  $Z^n(C)$ ,  $\varphi > 0$ . Suppose for any matrix  $C' \in \mathfrak{R}(\varphi)$  and  $x \in X(x^0)$  there exists an index  $i \in N_n$  such that

$$q_i(x, x^0, C_i + C'_i) > 0.$$

Then

$$x^0 \in P^n(C + C')$$

for any matrix  $C' \in \mathfrak{R}(\varphi)$ .

**Proof.** Let  $x \notin X(x^0)$ . Then for any  $t \in T$  there exists  $t' \in T$  such that  $N(x^0, t) = N(x, t')$ . Hence we have for any index  $i \in N_n$  and any matrix  $C' \in \mathfrak{R}(\varphi)$

$$\begin{aligned} q_i(x, x^0, C_i + C'_i) &= \max_{t \in T} (C_i + C'_i)[t]x - \max_{t \in T} (C_i + C'_i)[t]x^0 = \\ &= \max_{t \in T} (C_i + C'_i)[t]x - (C_i + C'_i)[t^*]x^0 \geq (C_i + C'_i)[t']x - (C_i + C'_i)[t^*]x^0 = 0. \end{aligned}$$

It means that

$$x^0 \in P^n(C + C')$$

for any matrix  $C' \in \mathfrak{R}(\varphi)$ . Lemma 2 is proved.

For any non-trivial solution  $x^0$  put

$$\varphi^n(x^0, C) = \min_{x \in X(x^0)} \max_{i \in N_n} \min_{t^0 \in T(x^0, x)} \max_{t \in T} \frac{C_i[t]x - C_i[t^0]x^0}{\sigma(x^0, t^0, x, t)},$$

where

$$\sigma(x^0, t^0, x, t) = |(N(x^0, t^0) \cup N(x, t)) \setminus (N(x^0, t^0) \cap N(x, t))|.$$

The following statements are true

$$t^0 \in \bar{T}(x, x^0) \implies \forall t \in T \ (\sigma(x^0, t^0, x, t) = 0). \quad (1)$$

$$C_i[t]x - C_i[t^0]x^0 + \|C_i\|_\infty \sigma(x^0, t^0, x, t) \geq 0, \quad i \in N_n, \quad (2)$$

It is easy to see that  $0 \leq \varphi^n(x^0, C) < \infty$ .

**Theorem 2** *The stability radius  $\rho^n(x^0, C)$  of any non-trivial efficient solution  $x^0$  of the problem  $Z^n(C)$  is expressed by the formula*

$$\rho^n(x^0, C) = \varphi^n(x^0, C).$$

**Proof.** First let us prove that  $\rho^n(x^0, C) \geq \varphi := \varphi^n(x^0, C)$ . For  $\varphi = 0$ , it is nothing to prove. Let  $\varphi > 0$ . Then for any  $x \in X(x^0)$  there exists an index  $i \in N_n$  such that

$$\min_{t^0 \in T(x^0, x)} \max_{t \in T} \frac{C_i[t]x - C_i[t^0]x^0}{\sigma(x^0, t^0, x, t)} \geq \varphi.$$

We have the following statements for any  $C' \in \mathfrak{R}(\varphi)$

$$\begin{aligned} q_i(x, x^0, C_i + C'_i) &= \max_{t^0 \in T} (C_i + C'_i)[t]x - \max_{t^0 \in T} (C_i + C'_i)[t^0]x^0 = \\ &= \min_{t^0 \in T} \max_{t \in T} (C_i[t]x - C_i[t^0]x^0 + C'_i[t]x - C'_i[t^0]x^0) \geq \\ &\geq \min_{t^0 \in T} \max_{t \in T} (C_i[t]x - C_i[t^0]x^0 - \|C'_i\| \sigma(x^0, t^0, x, t)). \end{aligned}$$

Using (1) we continue

$$= \min_{t^0 \in T(x^0, x)} \max_{t \in T} (C_i[t]x - C_i[t^0]x^0 - \|C'_i\| \sigma(x^0, t^0, x, t))$$

Applying (2) we finally conclude

$$> \min_{t^0 \in T(x^0, x)} \max_{t \in T} (C_i[t]x - C_i[t^0]x^0 - \varphi \sigma(x^0, t^0, x, t)) \geq 0.$$

Thus, by lemma 2, we obtain that non-trivial solution  $x^0$  preserves efficiency for any perturbing matrix  $C' \in \mathfrak{R}(\varphi)$ , i.e.  $\rho^n(x^0, C) \geq \varphi$ .

It remains to check that  $\rho^n(x^0, C) \leq \varphi$ . According to the definition of  $\varphi$ , there exists  $x \in X(x^0)$  such that for any  $i \in N_n$

$$\varphi \geq \min_{t^0 \in T(x^0, x)} \max_{t \in T} \frac{C_i[t]x - C_i[t^0]x^0}{\sigma(x^0, t^0, x, t)} = \max_{t \in T} \frac{C_i[t]x - C_i[\tilde{t}]x^0}{\sigma(x^0, \tilde{t}, x, t)}. \quad (3)$$

Let  $\varepsilon > 0$ . Consider the following perturbing matrix  $C^* \in \mathbf{R}^{n \times m}$ . Every string  $C_i^*$ ,  $i \in N_n$  of this matrix consists of the elements

$$C_{ij}^* = \begin{cases} \alpha & \text{if } j \in N(x^0, \tilde{t}), \\ -\alpha & \text{otherwise,} \end{cases}$$

where  $\varphi < \alpha < \varepsilon$ . Using (3) we get the following expressions:

$$\begin{aligned} q_i(x, x^0, C_i + C_i^*) &= \max_{t \in T} (C_i + C_i^*)[t]x - \max_{t \in T} (C_i + C_i^*)[t]x^0 \leq \\ &\max_{t \in T} (C_i + C_i^*)[t]x - (C_i + C_i^*)[\tilde{t}]x^0 = (C_i + C_i^*)[\hat{t}]x - (C_i + C_i^*)[\tilde{t}]x^0 = \\ &= C_i[\hat{t}]x - C_i[\tilde{t}]x^0 - \alpha \sigma(x^0, \tilde{t}, x, \hat{t}) < C_i[\hat{t}]x - C_i[\tilde{t}]x^0 - \varphi \sigma(x^0, \tilde{t}, x, \hat{t}) \leq \end{aligned}$$

$$\begin{aligned} &\leq C_i[\hat{t}]x - C_i[\tilde{t}]x^0 - \sigma(x^0, \tilde{t}, x, \hat{t}) \max_{t \in T} \frac{C_i[t]x - C_i[\tilde{t}]x^0}{\sigma(x^0, \tilde{t}, x, t)} \leq \\ &\leq C_i[\hat{t}]x - C_i[\tilde{t}]x^0 - \sigma(x^0, \tilde{t}, x, \hat{t}) \frac{C_i[\hat{t}]x - C_i[\tilde{t}]x^0}{\sigma(x^0, \tilde{t}, x, \hat{t})} = 0. \end{aligned}$$

Hence  $x^0$  is not efficient solution of the problem  $Z^n(C + C^*)$ , where  $C^* \in \mathfrak{R}(\varphi)$ . It means that  $\rho^n(x^0, C) \leq \varphi$ . This completes the proof of Theorem 2.

Assume that  $T = \{t^5\}$ .  $t^0 = \begin{bmatrix} 1 & 2 & \dots & m \\ 1 & 2 & \dots & m \end{bmatrix}$ . Then our problem transforms into vector linear Boolean programming problem

$$f_i(x, C_i) = C_i x \longrightarrow \min_{x \in X}, \quad i \in N_n,$$

where  $X \subseteq \mathbf{E}^m$ .

In this case one can see that any efficient solution is non-trivial. The next corollary follows from theorem 2.

**Corollary 1** [1] *The stability radius of any efficient solution  $x^0$  of vector linear Boolean programming problem  $Z^n(C)$ ,  $n \geq 1$ , equals to*

$$\min_{x \in X \setminus \{x^0\}} \max_{i \in N_n} \frac{C_i(x - x^0)}{\|x - x^0\|^*},$$

where  $\|z\|^* = \sum_{j \in N_n} |z_j|$ ,  $z = (z_1, z_2, \dots, z_m) \in \mathbf{R}^m$ .

Any efficient solution  $x^0$  of the problem  $Z^n(C)$  is called *stable* if  $\rho^n(x^0, C) > 0$ , and *strongly efficient* if there does not exist  $x \in X \setminus \{x^0\}$  such that  $C_i x^0 \geq C_i x$ . From corollary 2 we have

**Corollary 2** [1] *Any efficient solution of vector linear Boolean programming problem is stable iff it is strongly efficient.*

## References

- 1] Emelichev V.A., Girlich E., Podkopaev D.P. On stability of efficient solution in a vector trajectorial problem of discrete optimization, I,II, *Izv. Akad. Nauk Resp. Moldova. Matem.*, 1996, No. 3, pp. 5 – 16; 1997, No. 2, pp. 9 – 25. (Russian)
- [2] Emelichev V.A., Krichko V.N. On stability of Smale optimal, Pareto optimal and Slater optimal solution in vector trajectorial optimization problems, *Izv. Akad. Nauk Resp. Moldova. Matem.*, 1998, No. 3, pp. 81 – 86.
- [3] Emelichev V.A., Stepanishina Yu. V. Stability of majority efficient solution of a vector linear trajectorial problem, *Computer Science Journal of Moldova*, 1999, vol. 7, No. 3, pp. 291 – 306.
- [4] Emelichev V.A., Nikulin Yu.V. On the stability of efficient solution in a vector quadratic Boolean programming problem, *Izv. Akad. Nauk Resp. Moldova. Matem.*, 2000, No. 1, pp. 33 – 40.

V.A.Emelichev, V.N.Krichko, Yu.V.Nikulin,

Received April 30, 2002

Belarusian State University  
a. Fr. Skoryna, 4,  
Minsk 220050, Belarus  
E-mail: *emelichev@bsu.by*