

BeyondBenford R Package to compare Benford's and BDS's Distributions

Stéphane Blondeau Da Silva

Abstract

The package **BeyondBenford** compares the goodness of fit of Benford's and Blondeau Da Silva's (BDS's) digit distributions in a given dataset. It first enables to check whether the data distribution is consistent with theoretical distributions highlighted by Blondeau Da Silva or not; indeed, this ideal theoretical distribution must be at least approximately followed by the data for the use of BDS's model to be well-founded. It also allows to draw histograms of digit frequencies or probabilities (and their confidence or prediction intervals), both observed in the dataset and given by the two theoretical approaches. Finally, it proposes to quantify the goodness of fit of these laws via Pearson's chi-squared tests.

Keywords: Benford's law, digits, experimental data, R-package.

MSC 2010: 60E05.

1 Introduction

Benford's Law, also called Newcomb-Benford's Law, is somewhat surprising; indeed, the first digit d , $d \in \llbracket 1, 9 \rrbracket$, of numbers in many naturally occurring collections of data does not follow a discrete uniform distribution, as might be thought, but a logarithmic distribution (see the recent books of [1] and [2]). Discovered by the astronomer Newcomb in 1881 [3], it was definitively brought to light by the physicist Benford in 1938 [4]. The probability that d is the first digit of a number

is approximately:

$$\log\left(1 + \frac{1}{d}\right) .$$

It can also be extended to digits beyond the first one [5], the probability for d , $d \in \llbracket 0, 9 \rrbracket$, to be the p^{th} digit of a number being:

$$\sum_{j=10^{p-2}}^{10^{p-1}-1} \log\left(1 + \frac{1}{10j + d}\right) .$$

It was quickly admitted that numerous empirical data sets follow Benford's law: economic data [6], social data [7], demographic data [8], [9], physical data [10], [11] or biological data [12], [13] for instance; to such an extent that this law was used to detect possible frauds in lists of socio-economic data [14], [15] or in scientific publications [16].

Nevertheless many discordant voices brought a significantly different message. By putting aside the distributions known to fully disobey Benford's law [17]–[19], this law often appeared to be a good approximation of the reality, but no more than an approximation [20]–[22].

Similar to first digit case, the distributions of digits beyond the first have been observed in various application areas [11], [16], [23] and have also been used to detect frauds [24]–[26]. Once more, limits of such methods were emphasized [24], [25], [27].

Blondeau Da Silva, considering data as realizations of a homogeneous and expanded range of random variables following discrete uniform distributions, showed that, the proportion of each d , $d \in \llbracket 0, 9 \rrbracket$, as leading digit [28] or as other digit [29] structurally fluctuates. He demonstrated that, in his models, the predominance of 1 as digit (followed by 2 and so on) is all but surprising, and that the observed fluctuations around the values of probability determined by Benford's Law are also predictable: there is not a single Benford's Law but numerous distinct laws each of them determined by a parameter, the upper-bound of the considered data.

The huge and growing literature on Benford's law is available on the online database www.benfordonline.net with well over 1000 papers.

Two R packages [30] already enable to check whether datasets conform to Benford's law or not: **BenfordTests** [31] and **benford.analysis** [32]. The package **BeyondBenford** [33] compares the goodness of fit of Benford's law, on the one hand, and BDS's laws, on the other hand. Indeed, these latter, under certain conditions that we will recall, allow a better reliability of adjustment. The package **BeyondBenford** calculates the digit distribution in the considered dataset and determines whether it is consistent with BDS's or Benford's one. It also provides plotting tools for the visual evaluation of these distributions. We will walk through a detailed example to give an overview of the **BeyondBenford** package.

2 An example to get familiar with the main functions

Street addresses of Pierre-Buffiere, a small town of approximately 1200 inhabitants in Haute-Vienne (France), are available on:

www.data.gouv.fr/fr/datasets/base-d-adresses-nationale-ouverte-bano/,

which is an open platform for French public data.

After loading the package (`library(BeyondBenford)`), the code `data(address_PierreBuffiere)` gives us access to the sample data. This factor contains 346 rows, with each row representing an address number.

3 Is the data consistent with BDS's model?

This is the first essential question that must be answered in the affirmative. If this is not the case, comparisons are not relevant: the package should not be used. Indeed the use of the package is appropriate when the studied data can be considered as realizations of a homogeneous and expanded range of random variables approximately following discrete uniform distributions. In this model, the data is strictly positive and is lower and upper-bounded, constraint which is often valid in datasets,

the physical, biological, demographic, social and economical quantities being limited [28].

Among the different domains studied by Benford [4], some could be well adapted to our model: sizes of populations or street addresses for example (see [28] for a detailed explanation). [34] advised precisely to use their own similar model in the case of street addresses or when considering the first-page numbers of articles in a bibliography.

[29] showed that the model induces a specific distribution of positive integers determined by a lower and an upper-bound. Hence, in order to conform as closely as possible to the model, the studied database must have a distribution similar to that described in [29]. In p^{th} digit case, the probability p_k to obtain the number $k \in \llbracket l_b; u_b \rrbracket$ (where l_b is the lower-bound and u_b is the upper-bound) verifies:

$$p_k = \frac{1}{u_b - l_b + 1} \sum_{i=k-l_b+1}^{u_b-l_b+1} \frac{1}{i} .$$

In the studied example, the minimum value of the street number is 1 and the maximum value is 74. The associated theoretical distribution for the second digit is plotted in Figure 1.

The **BeyondBenford** package provides plotting tools to determine whether the data is consistent with BDS's model: the function `dat.distr`. The function's main arguments are as follows:

- **dat**: the considered dataset, a data frame containing non-zero real numbers.
- **theor**: if `theor=TRUE` BDS's theoretical distribution is plotted, otherwise only the histogram is represented. Default value: `theor=TRUE`.
- **nclass**: a strictly positive integer: the number of classes in the histogram. Default value: `nclass=50`.
- **conv**: if `conv=1`, all values of the dataset are multiplied by 10^k where k is the smallest positive integer such that all non-zero




Figure 1. Theoretical distribution of street numbers in the second digit case, the upper-bound being 74.

numerical values in the newly multiplied data frame have an absolute value greater than or equal to 1. Default value: `conv=0`.

- **lbound**: a positive integer, which characterizes the data. All (or most) of the data are greater than this "lower-bound". Default value: `lbound=max(floor(min(abs(dat)))+1,(10**(dig-1)))`.
- **upbound**: a positive integer, which characterizes the data. All (or most) of the data are lower than this "upper-bound". Default value: `upbound=ceiling(max(dat))`.
- **dig**: the chosen position of the digit (from the left). Default value: `dig=1`.
- **nchi**: the number of classes for values from 10^{p-1} to $\max(\max(\text{dat}), \text{upbound})$. If `nchi>0`, the function returns the chi-squared statistic (with `nchi - 1` degrees of freedom) of goodness of fit determined by the different classes. The null hypothesis states

that the studied distribution is consistent with the considered theoretical distribution. Default value: `nchi=0`.

Let us apply the `dat.distr` function to the `address_PierreBuffiere` dataset. The output from `dat.distr` is the data histogram along with optional BDS's theoretical distributions (Figure 2).

It can be noted that there are also other arguments for the formatting of histograms.

Example 1. *## Both the histogram and theoretical distribution are represented*

```
dat.distr(address_PierreBuffiere, dig=2, nclass=65)
```

Note that, out of the 346 values, only 217 are taken into consideration here because the values need to have at least 2 digits.

The data distribution looks similar to the one described by BDS's model (Figure 2). Let us provide a second example in which we numerically determine whether the studied distribution is conform to the theoretical distribution or not:

Example 2. *## The function returns the chi-squared statistic of goodness of fit determined by nchi classes.*

```
dat.distr(address_PierreBuffiere, dig=2, nchi=4)
```

```
[1] "Class freq.:" "130" "51" "25"
[5] "11"
[1] "Theor. freq.:" "127.050977504473" "54.6446927640259"
"26.7009825219254"
[5] "8.60334720957567"
```

```

                chi2                pval
1 Chi2 value is: The p-value is:
2 1.08754607580421 0.780081329402347
```

The `dat.distr` function returns, if requested, the frequencies of each equal-sized class of the dataset and the associated theoretical frequencies. It also returns a data frame containing the value of the chi squared test-statistic and its p-value. Note that the number of classes is limited by the theoretical frequencies that cannot exceed 5 in Pearson's chi-squared test [35]. In our example, the null hypothesis cannot




Figure 2. The data histogram of street numbers in the second digit case and its BDS's theoretical distribution.

be rejected: the studied distribution is consistent with the theoretical distribution.

4 Comparisons of the goodness of fit of Benford's and BDS's digit distributions in a dataset

4.1 Raw data

The package **BeyondBenford** provides a function returning the frequencies of each figure at a given position in the considered dataset: `obs.numb.dig`. Its two arguments `dat` and `dig` have already been defined in `dat.distr` function. The function output is a vector containing the frequencies of each figure in ascending order.

Let us give an example:

Example 3. `obs.numb.dig(address_PierreBuffiere, dig=2)`
[1] 31 24 27 21 25 17 21 16 19 16

For instance, there are 25 values with the second digit being a 4.

The package **BeyondBenford** also provides a function returning Benford's probability that a figure is at a given position: `Benf.val`. Its argument `dig` has already been defined in `dat.distr` function. Its second argument `fig` stands for the considered figure.

Let us give an example:

Example 4. `Benf.val(4, dig=2)`
[1] 0.1003082

The package **BeyondBenford** at last provides a function returning BDS's probability that a figure is at a given position (once the associated upper-bound has been specified): `Blon.val`. Its four arguments – `dig`, `lbound`, `ubound`, and `fig` – have already been defined above.

Let us give an example:

Example 5. `Blon.val(fig=4, dig=2, ubound=74)`
[1] 0.09836642

In Table 1 below, the frequencies of each figure in the considered dataset, regarding the second digit of address numbers, are listed.

Table 1. Values of frequency of each figure as second digit in the database, Benford's and BDS's theoretical values (the chosen upper-bound being 74). These values are rounded to the nearest tenthousandth.

Figure	Freq. in the database	BDS's values	Benford's values
0	0.1429	0.1436	0.1197
1	0.1106	0.1247	0.1139
2	0.1244	0.1136	0.1088
3	0.0968	0.1053	0.1043
4	0.1152	0.0984	0.1003
5	0.0783	0.0924	0.0967
6	0.0968	0.0872	0.0934
7	0.0737	0.0825	0.0904
8	0.0876	0.0782	0.0876
9	0.0737	0.0742	0.0850

The BDS's theoretical values seem slightly better; in particular the frequency range is higher, both for the observed data and for BDS's theoretical values.

4.2 Plotting tools

The **BeyondBenford** package provides plotting tools to perform the comparison between the two models with the function `digit.distr`. In addition to arguments that are shared with `dat.distr` (**dat**, **dig**, **lbound**, **upbound**), the `digit.distr` function has mainly the following additional arguments:

- **mod**: if `mod="ben"`, the data histogram and that of Benford are displayed, if `mod="ben&b1o"`, the data histogram, that of Benford and that of BDS are plotted, and otherwise the data histogram and that of BDS are given. Default value: `mod="ben"`.

- **No.sd**: the positive decimal number of standard deviation that defines the confidence and prediction intervals i.e. the error bars. If No.sd=0, no error bars are drawn. Default value: No.sd=0.
- **Sd.pr**: If Sd.pr=1, error bars for proportions are plotted (with No.sd standard deviation confidence intervals). If Sd.pr=0, they are not plotted. Default value: Sd.pr=0.

Let us apply the `digit.distr` function to the `address_PierreBuffiere` dataset. The output from `dat.distr` is a histogram of theoretical and experimental digit distribution (Figure 3).

There are also other arguments for the formatting of histograms.

Example 6. `digit.distr(address_PierreBuffiere, dig=2, mod="ben&blo", No.sd=1, Sd.pr=1)`

Naturally, Figure 3 is consistent with Table 1.

4.3 Pearson's chi-squared test

To quantify the quality of theoretical models, we use Pearson's chi-squared test of goodness of fit [35]: the null hypothesis states that the studied distribution is consistent with the considered theoretical distribution, *i.e.* Benford's or BDS's ones. The function `chi2` determines the test statistic and its associated p-value. In addition to arguments that are shared with `dat.distr` (`dat`, `dig`, `lbound`, `ubound`), the `chi2` function has the following specific arguments:

- **mod**: if mod="ben", the theoretical distribution considered is that of Benford, else it is BDS's ones which is chosen. Default value: mod="ben".
- **pval**: if pval=0, the p-value is not returned, else it is available. Default value: pval=0.

Let us apply this function to the `address_PierreBuffiere` dataset. The output from `chi2` is a data frame containing the Pearson's chi-squared statistic (and the associated p-value if requested).




Figure 3. The histogram of Pierre Buffière’s street numbers in the second digit case along with Benford’s and BDS’s distributions of digit.

Example 7. *## Measure of Benford's Law goodness of fit*

```
chi2(address_PierreBuffiere, dig=2, pval=1)
```

```
      chi2      pval
```

```
1 Chi2 value is: The p-value is:
```

```
2 3.84793221030181 0.921132993758269
```

Measure of BDS's Law goodness of fit

```
chi2(address_PierreBuffiere, dig=2, pval=1, mod="BDS")
```

```
      chi2      pval
```

```
1 Chi2 value is: The p-value is:
```

```
2 2.47996278328848 0.981417506807756
```

In both cases the null hypothesis cannot be rejected: the studied distribution is consistent with the theoretical distributions. It can be noted that the quality of the adjustment seems slightly better with BDS's model.

5 Conclusion

The use of Benford's Law has increased rapidly in the last few years in extremely diverse fields such as mathematics, physics, biology, economics and demography, to name but a few. But the adjustment proposed by Benford is often only approximate. In some precisely described cases, BDS's probability distributions should be preferred. Indeed probabilities of occurrence of digits in these distributions fluctuate around Benford's values [28], [29].

The **BeyondBenford** package is thus a relevant tool to compare the goodness of fit of Benford's and BDS's distributions in a given collection of data. It offers new laws to find a better approximation of digits distribution in a considered dataset.

References

- [1] S. J. Miller, Ed., *Benford's Law: Theory and Applications*. Princeton, NJ: Princeton University Press, 2015.

- [2] A. Berger and T. Hill, *An Introduction to Benford's Law*. Princeton, NJ: Princeton University Press, 2015.
- [3] R. Newcomb, "Note on the frequency of use of the different digits in natural numbers," *American Journal of Mathematics*, vol. 4, pp. 39–40, 1881.
- [4] F. Benford, "The law of anomalous numbers," *Proceedings of the American Philosophical Society*, vol. 78, pp. 127–131, 1938.
- [5] T. Hill, "The significant-digit phenomenon," *The American Mathematical Monthly*, vol. 102, no. 4, pp. 322–327, 1995.
- [6] T. Sehity, E. Hoelz, and E. Kirchler, "Price developments after a nominal shock: Benford's Law and psychological pricing after the euro introduction," *International Journal of Research in Marketing*, vol. 22, no. 4, pp. 471–480, 2005.
- [7] J. Golbeck, "Benford's law applies to online social networks," *Plos One*, vol. 10, no. 8, 2015.
- [8] M. Nigrini and W. Wood, "Assessing the integrity of tabulated demographic data," 1995, preprint.
- [9] L. Leemis, B. Schmeiser, and D. Evans, "Survival distributions satisfying Benford's Law," *The American Statistician*, vol. 54, no. 4, pp. 236–241, 2000.
- [10] M. Nigrini and S. Miller, "Benford's Law applied to hydrology data—results and relevance to other geophysical data," *Mathematical Geology*, vol. 39, no. 5, pp. 469–490, 2007.
- [11] T. Alexopoulos and S. Leontsinis, "Benford's Law in astronomy," *Journal of Astrophysics and Astronomy*, vol. 35, no. 4, pp. 639–648, 2014.
- [12] E. Costasa, V. Lopez-Rodasa, F. Torob, and A. Flores-Moya, "The number of cells in colonies of the cyanobacterium *microcystis aeruginosa* satisfies benford's law," *Aquatic Botany*, vol. 89, no. 3, pp. 341–343, 2008.

- [13] J. L. Friar, T. Goldman, and J. Pérez-Mercader, “Genome sizes and the Benford distribution,” *Plos One*, vol. 7, no. 5, 2012.
- [14] C. Durtschi, W. Hillison, and C. Pacini, “The effective use of benford’s law to assist in detecting fraud in accounting data,” *Journal of Forensic Accounting*, vol. V, pp. 17–34, 2004.
- [15] K. Tödter, “Benford’s Law as an indicator of fraud in economics,” *German Economic Review*, vol. 10, pp. 339–351, 2009.
- [16] A. D. Alves, H. H. Yanasse, and N. Y. Soma, “Benford’s Law and articles of scientific journals: comparison of JCR and Scopus data,” *Scientometrics*, vol. 98, pp. 173–184, 2014.
- [17] R. A. Raimi, “The first digit problem,” *American Mathematical Monthly*, vol. 83, no. 7, pp. 521–538, 1976.
- [18] T. Hill, “Random-number guessing and the first digit phenomenon,” *Psychological Reports*, vol. 62, no. 3, pp. 967–971, 1988.
- [19] T. W. Beer, “Terminal digit preference: beware of benford’s law,” *Journal of Clinical Pathology*, vol. 62, no. 2, p. 192, 2009.
- [20] P. D. Scott and M. Fasli, “Benford’s Law: an empirical investigation and a novel explanation,” CSM Technical Report 349, University of Essex, 2001, <https://cswww.essex.ac.uk/technical-reports/2001/CSM-349.pdf>.
- [21] A. Saville, “Using Benford’s Law to detect data error and fraud: An examination of companies listed on the Johannesburg Stock Exchange,” *South African Journal of Economic and Management Sciences*, vol. 9, no. 3, pp. 341–354, 2006.
- [22] N. Gauvrit and J.-P. Delahaye, *Scatter and Regularity Imply Benford’s Law... and More*. World Scientific, 2011, pp. 53–69.
- [23] D. Geyer, “Detecting fraud in financial data sets,” *Journal of Business and Economics Research*, vol. 8, no. 7, pp. 75–83, 2010.

- [24] W. R. Mebane Jr, “Election forensics: The second-digit benford’s law test and recent american presidential elections,” in *Proceedings of the Election Fraud Conference*, 2006.
- [25] W. K. T. Cho and B. J. Gaines, “Breaking the (Benford) Law: Statistical fraud detection in campaign finance,” *The American Statistician*, vol. 61, no. 3, pp. 218–223, 2007.
- [26] D. W. Joenssen, “Two digit testing for benford’s law,” in *Proceedings of the ISI World Statistics Congress, 59th Session in Hong Kong*, 2013.
- [27] A. Diekmann, “Not the first digit! Using benford’s law to detect fraudulent scientific data,” *Journal of Applied Statistics*, vol. 34, no. 3, pp. 321–329, 2007.
- [28] S. Blondeau Da Silva, “Benford or not Benford: a systematic but not always well-founded use of an elegant law in experimental fields,” *Communications in Mathematics and Statistics*, 2019.
- [29] —, “Benford or not Benford: new results on digits beyond the first,” *arXiv:1805.01291 [stat.OT]*, 2018.
- [30] R Core Team, *R: A Language and Environment for Statistical Computing*, R Foundation for Statistical Computing, Vienna, Austria, 2013. [Online]. Available: <http://www.R-project.org/>
- [31] D. W. Joenssen, *BenfordTests: Statistical Tests for Evaluating Conformity to Benford’s Law*, 2015, r package version 1.2.0. [Online]. Available: <https://CRAN.R-project.org/package=BenfordTest>
- [32] C. Cinelli, *benford.analysis: Benford Analysis for Data Validation and Forensic Analytics*, 2018, r package version 0.1.5. [Online]. Available: <https://CRAN.R-project.org/package=benford.analysis>
- [33] S. Blondeau Da Silva, *BeyondBenford: Compare the Goodness of Fit of Benford’s and Blondeau Da Silva’s Digit Distributions to a*

Given Dataset, 2020, r package version 1.3. [Online]. Available: <https://CRAN.R-project.org/package=BeyondBenford>

- [34] E. Janvresse and T. De La Rue, “From uniform distributions to Benford’s Law,” *Journal of Applied Probability*, vol. 41, no. 4, pp. 1203–1210, 2004.
- [35] K. Pearson, “On the criterion that a given system of deviations from the probable in the case of a correlated system of variables is such that it can be reasonably supposed to have arisen from random sampling,” *Philosophical Magazine*, vol. 50, no. 302, pp. 157–175, 1900.

Stéphane Blondeau Da Silva,

Received August 30, 2020

Accepted July 25, 2021

Stéphane Blondeau Da Silva
Institution Académie de Limoges
Address 13 Rue François Chénieux, 87000 Limoges France
Phone: 07 81 49 13 18
E-mail: Stephane.Blondeau-Da-Silva@ac-limoges.fr